

CASE REPORT

Open Access



Isolated sternal fracture after low-energy trauma in a geriatric patient: a case report

Joni J. Sairanen¹ and Otso Arponen^{2,3*}

Abstract

Background: Falls are a common cause of emergency department (ED) visits for the older population. If osteoporosis is present, even falls from standing height can lead to unusual fractures normally associated with high-energy trauma. In this report, we analyze a rare case of an isolated sternal fracture with an unusual mechanism of injury. Our discussion aims to improve care for older adults with fall-related fractures.

Case presentation: An 86-year-old female presented in the ED of our hospital with a complaint of chest pain. She recalled a fall at home the previous day and described how her fist was impacted between the floor and her chest. A physical examination revealed local tenderness in the mid-chest. A lateral chest x-ray indicated a sternal fracture, and a chest computed tomography scan ruled out concomitant injuries. There were no acute changes on her electrocardiogram (ECG). Conservative outpatient treatment was started, and referrals were made with a recommendation to initiate fall prevention measures and osteoporosis screening in primary health care.

Conclusions: Geriatric patients can present in the ED with a rare sternal fracture even after only a minor chest trauma. Appropriate imaging and an ECG are warranted to exclude life-threatening additional injuries. An in-depth physical examination and an understanding of the exact mechanism of injury are important to avoid missing fractures in unexpected locations. Modern ED physicians could have an important role in the secondary prevention of fall-related fractures for geriatric patients.

Keywords: Sternal fracture, Trauma imaging, Falls, Osteoporosis, Case report

Background

Of all trauma admissions in the emergency department (ED), only 0.33–2.1% involve a sternal fracture (SF) due to blunt chest trauma [1, 2]. Motor vehicle accidents account for 83–84% of all SFs, while only 10–13% of SFs are due to falls, usually from height [3, 4]. Direct chest trauma is not the only possible mechanism of injury (MOI); a fall on the back could cause an SF due to flexion-compression [5]. Additional skeletal or visceral injuries are present in approximately 80% of patients with an SF due to blunt chest trauma [1, 2]. An isolated SF is

diagnosed in the absence of concomitant injuries, and it can be described as a relatively mild injury [3, 6].

Eighty-seven percent of all fractures in persons older than 65 years are caused by falls [7]. The risk of fractures is markedly increased in patients with osteoporosis, which is both underdiagnosed and undertreated at the population level [8]. Potential risk factors for future falls and fractures should be identified and managed appropriately [7]. Even when a patient is discharged from the ED to go home, ED physicians could initiate secondary prevention by pinpointing the most obvious problems (e.g., inappropriate medication, poor vision or balance, orthostatic hypotension, frailty) and referring the patient to primary health care and osteoporosis screening.

*Correspondence: otso.arponen@tuni.fi

³ Faculty of Medicine and Health Technology, Tampere University, Tampere, Finland

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Case presentation

An 86-year-old female who lived alone at home arrived in the ED of our hospital and complained of chest pain that worsened with inspiration. She stated that the pain had started after a fall at home. Her medical records revealed a history of mild Alzheimer's disease and hypertension. A photocopy of a recent ambulance report was found, according to which paramedics had responded to a chest pain call at her apartment the previous day. A rib fracture had been suspected and the patient was advised to seek medical attention in primary health care, but she arrived in the hospital ED instead.

The patient was in no apparent pain or distress. She was able to walk unassisted, although the overall impression suggested frailty. Respiration rate was 17/min, oxygen saturation 97% on room air, pulse 58/min, and blood pressure 156/72 mmHg. Troponin T concentration was within normal limits, and hemoglobin level (13.7 g/dL) was normal. An electrocardiogram (ECG) showed sinus rhythm with a prolonged PR interval (.218 s) and a bifascicular block, which were also present on her previous ECGs and appeared unchanged.

A physical examination revealed local pain in the sternum, but there were no external signs of trauma to the patient's head, neck, or upper body. The rib cage was stable and nontender. Lung sounds were normal on auscultation. A systolic grade 2 murmur was noted. The cervical, thoracic, and lumbar spine were stable and nontender. No focal neurological deficits were noted.

The patient underwent imaging (Fig. 1), and the initial radiology report of the chest x-ray (CXR) came back as normal. The CXR was reviewed in the ED. On the lateral view (A), a cortical breach in the body of the sternum was noted. There were no signs of newly displaced rib fractures, pneumothorax, or hemothorax. A computed

tomography scan of the chest (B) confirmed the diagnosis of an isolated SF. No traumatic damage to major vessels or the heart could be discerned, but the ascending aorta was dilated (45 mm). Cortical thinning and loss of bony trabeculae were present, suggestive of osteopenia.

Upon further questioning, the patient gave a more precise description of the fall. She recalled how she fell forward, trapping her right fist between the floor and her chest. We believe this was an accurate account of the incident, despite the patient's problems with short-term memory. The MOI is unusual but plausible. The patient denied having had episodes of syncope or transient loss of consciousness, and she did not use alcohol or any drugs that could have contributed to the fall.

Conservative treatment was started for isolated SF. The exact cause of the fall remained obscure, but frailty was interpreted as a risk factor for recurrent falls and fractures in the future. The patient's relatives were informed, and the patient was discharged with prescriptions for acetaminophen and a vitamin D and calcium supplement. Written instructions were given to book a follow-up appointment with a primary care physician who could perform an osteoporosis workup and a comprehensive geriatric evaluation with the aim of preventing future falls. A cardiology referral was made due to the conduction abnormalities on the ECG, the systolic murmur, and the dilated ascending aorta.

Discussion and conclusions

The thoracic cavity houses many of our vital organs and vessels. Significant chest trauma indicates imaging to diagnose or rule out not only skeletal fractures but also severe concomitant injuries (e.g., pulmonary contusion, pneumothorax, and mediastinal hematoma) [1, 6, 9]. The possibility of cardiac contusion must be kept




Fig. 1 Isolated sternal fracture in a lateral chest x-ray (A) and a sagittal chest computed tomography view (B)

in mind: patients with new arrhythmias, hypotension, or ST-segment abnormalities should be monitored in the hospital [10]. Patients with isolated SF very rarely require surgical treatment, and their prognosis is generally favorable [11]. Based on level II evidence, they can be safely discharged home if there are no other indications for hospitalization (e.g., comorbidities, need for parenteral pain medication, or lack of social support) [4]. Conservative treatment of SF involves rest and analgesia [11]. Passive reduction and corset fixation could be needed in patients with displaced SF [11].

The SF of our patient was noticed only upon review of the CXR in the ED – possibly because the radiologist who compiled the initial radiology report did not have the complete clinical picture (the precise MOI was unknown at the time of the imaging request). It has been estimated that missed fractures account for up to 79.7% of all diagnostic errors in the ED [12]. Detailed history and all relevant clinical findings should be disclosed in imaging requests because they help in choosing the most appropriate imaging modality [13]. Better imaging requests could also reduce the possibility of radiograph misinterpretation [13]; this is particularly the case in blunt chest trauma, where the CXR has a poor screening performance with only 45% sensitivity and 78% negative predictive value for injury [14].

The role of ED in the treatment of fractures need not be limited to acute care. In a Danish study by Jantzen et al. [15], all ED patients older than 65 years with a low-energy Colles' fracture were systematically referred for osteoporosis screening. Sixty-five percent of the examined patients were diagnosed with osteoporosis, and almost all of them eventually received treatment with anti-osteoporosis drugs. Similarly, an ED physician could refer selected geriatric patients with any fall-related fractures to primary health care for comprehensive geriatric assessment and osteoporosis screening.

Fall-related fractures in older people are frequently encountered in the ED. Osteoporosis, a common and undertreated disease, predisposes patients to all types of fracture, including the rare SF. Radiologists' work could be made easier by disclosing detailed history and all relevant clinical findings in imaging requests. The modern ED physician could have an important role in the secondary prevention of fall-related fractures.

Abbreviations

ED: Emergency department; SF: Sternal fracture; MOI: Mechanism of injury; ECG: Electrocardiogram; CXR: Chest x-ray.

Acknowledgements

Not applicable.

Authors' contributions

JS and OA conceptualized the study, acquired the study data, and contributed to the writing and revision of the manuscript. JS interpreted the clinical data. OA prepared the images. All authors read and approved the final manuscript.

Funding

The authors received no funding for this study.

Availability of data and materials

All data generated or analyzed during this study are included in this published article.

Declarations

Ethics approval and consent to participate

Ethics approval is not applicable. The use of study data was approved by the Research Director of the Pirkanmaa Hospital District (R22622). The study was performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki and its later amendments. The authors have the patient's written informed consent for the participation in this study.

Consent for publication

The authors have the patient's written informed consent for the publication of this study and any accompanying images.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Emergency Department, Tampere University Hospital, Tampere, Finland.

²Department of Radiology, Tampere University Hospital, Tampere, Finland.

³Faculty of Medicine and Health Technology, Tampere University, Tampere, Finland.

Received: 6 June 2022 Accepted: 15 July 2022

Published online: 29 July 2022

References

- Recinos G, Inaba K, Dubose J, Barmparas G, Teixeira PG, Talving P, et al. Epidemiology of sternal fractures. *Am Surg*. 2009;75:401–4.
- Fokin AA, Wycech Knight J, Abid AT, Yoshinaga K, Alayon AL, Grady R, et al. Sternal fractures in blunt trauma patients. *Eur J Trauma Emerg Surg*. 2022. <https://doi.org/10.1007/s00068-021-01871-1>.
- von Garrel T, Ince A, Junge A, Schnabel M, Bahrs C. The sternal fracture: radiographic analysis of 200 fractures with special reference to concomitant injuries. *J Trauma*. 2004;57:837–44.
- Odell DD, Peleg K, Givon A, Radomislensky I, Makey I, DeCamp MM, et al. Sternal fracture: isolated lesion versus polytrauma from associated extrasternal injuries—analysis of 1,867 cases. *J Trauma Acute Care Surg*. 2013;75:448–52.
- Fowler AW. Flexion-compression injury of the sternum. *J Bone Joint Surg Br*. 1957;39-B:487–97.
- Harley DP, Mena I. Cardiac and vascular sequelae of sternal fractures. *J Trauma*. 1986;26:553–5.
- Ambrose AF, Cruz L, Paul G. Falls and fractures: a systematic approach to screening and prevention. *Maturitas*. 2015;82:85–93.
- Kanis JA, Svedbom A, Harvey N, McCloskey EV. The osteoporosis treatment gap. *J Bone Miner Res*. 2014;29:1926–8.
- Perez MR, Rodriguez RM, Baumann BM, Langdorf MI, Anglin D, Bradley RN, et al. Sternal fracture in the age of pan-scan. *Injury*. 2015;46:1324–7.
- Nagy KK, Krosner SM, Roberts RR, Joseph KT, Smith RF, Barrett J. Determining which patients require evaluation for blunt cardiac injury following blunt chest trauma. *World J Surg*. 2001;25:108–11.
- Klei DS, de Jong MB, Öner FC, Leenen LP, van Wessel KJ. Current treatment and outcomes of traumatic sternal fractures—a systematic review. *Int Orthop*. 2019;43:1455–64.
- Guly HR. Diagnostic errors in an accident and emergency department. *Emerg Med J*. 2001;18:263–9.

13. Hanna TN, Rohatgi S, Shekhani HN, Dave IA, Johnson JO. Clinical information available during emergency department imaging order entry and radiologist interpretation. *Emerg Radiol.* 2017;24:361–7.
14. Dillon DG, Rodriguez RM. Screening performance of the chest X-ray in adult blunt trauma evaluation: is it effective and what does it miss? *Am J Emerg Med.* 2021;49:310–4.
15. Jantzen C, Cieslak LK, Barzanji AF, Johansen PB, Rasmussen SW, Schmidt TA. Colles' fractures and osteoporosis—a new role for the emergency department. *Injury.* 2016;47:930–3.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

